




Edexcel A Level FM Revision Questions: Solutions

Polar curves

Solutions

1. (i) $(x - 1)^2 + y^2 = 1$



$$x = r\cos\theta \text{ and } y = r\sin\theta$$

$$\text{So } r^2\cos^2\theta + 1 - 2r\cos\theta + r^2\sin^2\theta = 1$$

$$\Rightarrow r^2 - 2r\cos\theta = 0$$

$$\Rightarrow r = 2\cos\theta \text{ or } r = 0$$

$$\text{ie } r = 2\cos\theta \text{ [with } r = 0 \text{ when } \theta = \frac{\pi}{2}\text{]}$$

$$\text{(ii) } r = \frac{2}{1+\cos\theta}; x = r\cos\theta \text{ and } y = r\sin\theta; \text{ also } r^2 = x^2 + y^2$$


$$\text{So } r + r\cos\theta = 2 \Rightarrow r = 2 - x \Rightarrow r^2 = (2 - x)^2$$

$$\Rightarrow x^2 + y^2 = 4 + x^2 - 4x \Rightarrow y^2 = 4(1 - x)$$

This can be obtained from the parabola $y^2 = 4x$ by the following steps:

$y^2 = 4(-x) = -4x$ [reflection in the y -axis; note that the curve now only exists for negative x]

$y^2 = -4(x - 1) = 4(1 - x)$ [translation of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$]




2.(i) $r = 5 + 4\cos\theta$

Step 1: As r is a function of $\cos\theta$, the curve will be symmetric about the x -axis.

Step 2: $r > 0$ at all times

Step 3: Key points to plot are $\theta = 0, r = 9$; $\theta = \frac{\pi}{2}, r = 5$; $\theta = \pi, r = 1$



(ii) The required x -coordinate can be found by investigating the vertical tangents; ie when $\frac{dx}{d\theta} = 0$ [when the x -coordinate is (instantaneously) not changing as θ changes]

$$x = r\cos\theta = (5 + 4\cos\theta)\cos\theta$$

$$\text{so that } \frac{dx}{d\theta} = (-4\sin\theta)\cos\theta + (5 + 4\cos\theta)(-\sin\theta) = -8\sin\theta\cos\theta - 5\sin\theta$$

$$\text{Then } \frac{dx}{d\theta} = 0 \Rightarrow \sin\theta = 0 \text{ (ie } \theta = 0 \text{ or } \pi) \text{ or } \cos\theta = -\frac{5}{8}$$

$$\Rightarrow x = (5 + 4\cos\theta)\cos\theta = \left(5 - \frac{20}{8}\right)\left(-\frac{5}{8}\right) = -\frac{25}{16}$$

$$\text{(iii) Area enclosed by curve} = 2 \int_0^{\pi} \frac{1}{2} (5 + 4\cos\theta)^2 d\theta$$

$$= \int_0^{\pi} 25 + 16\cos^2\theta + 40\cos\theta d\theta$$

$$= \int_0^{\pi} 25 + 8(1 + \cos 2\theta) + 40\cos\theta d\theta$$

$$= [33\theta + 4\sin 2\theta + 40\sin\theta]_0^{\pi}$$

$$= 33\pi$$

[Rough check: Area of rectangle of base 11 and height 10 is approx. 35π]


3. (i) Step 1: As $r = \pm\sqrt{\sin 2\theta}$ isn't a function of either $\cos\theta$ or $\sin\theta$, there is no symmetry about the x or y axis.

Step 2: The curve isn't defined for $\frac{\pi}{2} < \theta < \pi$ or for $\frac{3\pi}{2} < \theta < 2\pi$ (as $\sin 2\theta < 0$).

Step 3: For each θ there will positive and negative values of r of the same magnitude. [However the negative values of r for θ will overlap with the positive values for $\theta + \pi$.]

Step 4: Key points to plot are: $\theta = 0, r = 0$; $\theta = \frac{\pi}{4}, r = \pm 1$; $\theta = \frac{\pi}{2}, r = 0$ (and the cycle repeats itself for $\theta = \pi$ to $\theta = \frac{3\pi}{2}$).

Step 5: The gradient at $\theta = 0$ (when $r = 0$) is 0 (ie along the line $\theta = 0$), and at $\theta = \frac{\pi}{2}$ it is ∞ (ie along the line $\theta = \frac{\pi}{2}$).



(ii) $r = 1$ when $\theta = \frac{\pi}{4}$ for $r^2 = \sin 2\theta$, and when $\theta = 0$ for $r^2 = \cos 2\theta$, so the curve for $r^2 = \sin 2\theta$ needs to be rotated by $\frac{\pi}{4}$ clockwise.

[This rotation transforms $r^2 = \sin 2\theta$ to $r^2 = \sin 2(\theta + \frac{\pi}{4})$ [as clockwise is the negative direction] = $\sin(2\theta + \frac{\pi}{2}) = \cos 2\theta$]

$$(iii) y = r \sin \theta, r^2 = \sin 2\theta \Rightarrow y^2 = \sin 2\theta \cdot \sin^2 \theta$$

$$\Rightarrow 2y \frac{dy}{d\theta} = 2 \cos 2\theta \sin^2 \theta + \sin 2\theta (2 \sin \theta \cos \theta)$$

$$\text{Then } \frac{dy}{d\theta} = 0 \Rightarrow (\cos^2 \theta - \sin^2 \theta) \sin^2 \theta + (2 \sin \theta \cos \theta)(\sin \theta \cos \theta) = 0$$

$$\Rightarrow 3 \cos^2 \theta \sin^2 \theta - \sin^4 \theta = 0$$

$$\Rightarrow \sin^2 \theta (3 \cos^2 \theta - \sin^2 \theta) = 0$$

$$\Rightarrow \sin^2 \theta (3 \cos^2 \theta - (1 - \cos^2 \theta)) = 0$$

$$\Rightarrow \sin^2 \theta (4 \cos^2 \theta - 1) = 0$$

$$\Rightarrow \theta = 0 \text{ or } \pi \text{ (within } [0, 2\pi) \text{) (ie when the curve is at the Origin)}$$

$$\text{or } \cos \theta = \pm \frac{1}{2}, \text{ so that } \theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3} \text{ or } \frac{5\pi}{3}$$

The curve doesn't exist for $\theta = \frac{2\pi}{3}$ and $\frac{5\pi}{3}$, and so the required value is

$\theta = \frac{\pi}{3}$ (when the y -coordinate is positive).

$$\text{At } \theta = \frac{\pi}{3}, y^2 = \sin 2\theta \cdot \sin^2 \theta = \left(\frac{\sqrt{3}}{2}\right) \left(\frac{\sqrt{3}}{2}\right)^2 = \left(\frac{\sqrt{3}}{2}\right)^3$$

$$\Rightarrow y = \left(\frac{\sqrt{3}}{2}\right)^{\frac{3}{2}} = 0.80593 = 0.806 \text{ (3sf)}$$